



Chinese Journal of Wildlife
ISSN 2310-1490,CN 23-1587/S

2024-04-17
2024-12-23

[J/OL]

<https://link.cnki.net/urlid/23.1587.s.20241223.1015.002>



ISSN 2096-4188 CN 11-6037/Z

1#	1#	2	2
2	2	3	1*

1.

150040

2.

154200

Sika deer (*Cervus mandchuricus*) is an important ungulate animal in temperate forests, and also one of the main prey for the Amur tiger (*Panthera tigris*). The wild sika deer population of Heilongjiang Taipinggou National Nature Reserve, which serves as a cross-border corridor for the Amur tiger was extinct in the 1980s. It has been planned to promote the rapid recovery of the wild population of sika deer through the domestication of captive sika deer behavior within the reserve. However, the instantaneous scanning sampling method has been widely used in behavioral research, but there is relatively little scientific research on the specific sampling interval of instantaneous scanning. Therefore, the different behavioral durations of sika deer were obtained at different sampling intervals of 5 min and 10, 15, 20, 30, 60 min. The linear regression method was used to compare the different behavioral durations in pairs. The three criteria, $r^2 \geq 0.9$, intercept no significant difference from 0 ($P > 0.05$), and slope no significant difference from 1 ($P > 0.05$), were used to measure whether the sampling interval can accurately estimate the duration of the behavior, and then determine the optimal sampling interval. The results indicated that the optimal sampling interval for locomotive behavior was 5 min, the optimal sampling interval for

[14]


[15]

[16-17]

3 1 1 0.2 1

Period	/d Days	/ g · ⁻¹ Total amount of feed per day for each	/ d ⁻¹ Frequency of feed per day
Epidemic quarantine period	14 0~14	1 000	2

Rewilding training period



Behavior	Sample size	/h Duration of behavior					
		/d	5 min	10 min	15 min	20 min	30 min
			5	1	1	2	3
Resting	20		9.28 ± 2.49	9.27 ± 2.50	9.26 ± 2.33	9.32 ± 2.59	9.28 ± 2.24
Ingestive	20		7.58 ± 3.42	7.61 ± 3.48	7.56 ± 3.33	7.49 ± 3.53	7.63 ± 3.33

9.27 ± 2.43

7.52 ± 3.48

Ruminating	20	1.81 ± 0.85	1.79 ± 0.84	1.89 ± 0.81	1.81 ± 0.84	1.75 ± 0.68	1.88 ± 0.68
Locomotive	20	$1.61 \pm$					

7

7

7

[32]

7

10 min 15 min

[33]

10 min

15 min

5 min

2

0.9

[34]

5 min 5 min

[7,35-37]

10 min

5 min

15 min

[1] , , . 1 [M]. : , 2021:6. . 021
JIANG Z G, WU Y, LIU S Y, et al. China'

- [7] , , . [J]. , 2002, 21(6): 29-32.
LIU Z S, WU J P, TENG L W. Time budget and behavior pattern of sem-free in spring [J]. Chinese Journal of Ecology, 2002, 21(6): 29-32.
- [8] , , . [J]. , 2012, 40(1): 86-89.
LÜ S J, YANG Y, WEI W H. Activity rhythm of mature female sika deer in captive and semi-free conditions [J]. Journal of Northeast Forestry University, 2012, 40(1): 86-89.
- [9] BERNSTEIN I S. An empirical comparison of focal and ad libitum scoring with commentary on instantaneous scans, all occurrence and one-zero techniques [J]. Animal Behaviour, 1991, 42(5): 721-728.

- [21] ZHENG J L, CHENG K, LIU Y, et al. Effects of food and environmental enrichment on the diurnal behaviors of half free-living ring-tailed lemurs (*Lemur catta*) [J]. Chinese Journal of Wildlife, 2020, 41(2): 288-295.
- [22] WANG Y P, MA W H, SUN W. Diurnal behaviors time budgets and activity rhythm of captive snowleopards (*Uncia uncia*) in different age stages [J]. Chinese Qinghai Journal of Animal and Veterinary Sciences, 2023, 53(2): 1-5, 55. 2023 i c h m e n t
- [23] PULLIN A N, PAIRIS-GARCIA M D, CAMPBELL B J, et al. Technical note: Instantaneous sampling intervals validated from continuous video observation for behavioral recording of feedlot lambs [J]. Journal of Animal Science, 2017, 95(11): 4703-4707.
- [24] LEDGERWOOD D N, WINCKLER C, TUCKER C B. Evaluation of data loggers, sampling intervals, and editing techniques for measuring tk ues

LÜ S J, YANG L, YANG Y, et al. Research of diurnal activity rhythm of house-hold sika deer [J]. Ecological Science, 2009, 28(5): 433-

